arXiv:math/0701664v4 [math.GT] 12 Mar 2007

EXOTIC SMOOTH STRUCTURES ON SMALL 4-MANIFOLDS

ANAR AKHMEDOV AND B. DOUG PARK

Dedicated to Ronald J. Stern on the occasion of his sixtieth birthday

Abstract. Let M be either $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$ or $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$. We construct the first example of a simply-connected irreducible symplectic 4-manifold that is homeomorphic but not diffeomorphic to M.

1. Introduction

Given two 4-manifolds, X and Y, we denote their connected sum by $X\#Y$. For a positive integer $m \geq 2$, the connected sum of m copies of X will be denoted by mX for short. Let $\mathbb{CP}^2$ denote the complex projective plane and let $\overline{\mathbb{CP}}^2$ denote the underlying smooth 4-manifold $\mathbb{CP}^2$ equipped with the opposite orientation. There is an extensive and growing literature on the construction of exotic smooth structures on the closed 4-manifolds $\mathbb{CP}^2\#m\overline{\mathbb{CP}}^2$ and $3\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2$ for some small positive integers m and n. In the next two paragraphs, we briefly highlight some of the papers that have appeared.

The existence of an exotic smooth structure on a 4-manifold was first proved by Donaldson in [Do]. Using SU(2) gauge theory, he showed that a Dolgachev surface $E(1)_{2,3}$ is homeomorphic but not diffeomorphic to $\mathbb{CP}^2\#9\overline{\mathbb{CP}}^2$. Infinitely many irreducible smooth structures on $\mathbb{CP}^2\#9\overline{\mathbb{CP}}^2$ were later constructed by Fintushel, Friedman, Stern and Szabó in [Fri, FS2, Sz2]. Kotschick showed in [Ko1] that the Barlow surface in [Ba] is homeomorphic but not diffeomorphic to $\mathbb{CP}^2\#8\overline{\mathbb{CP}}^2$. More recently, J. Park was able to construct an exotic $\mathbb{CP}^2\#7\overline{\mathbb{CP}}^2$ in [Pa2] using the rational blowdown technique of Fintushel and Stern in [FS1]. Using a more general blowdown technique in [Pa1], Stipsicz and Szabó constructed an exotic $\mathbb{CP}^2\#6\overline{\mathbb{CP}}^2$ in [SS2]. In [FS3], Fintushel and Stern constructed infinitely many irreducible smooth structures on $\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2$ for $6 \leq n \leq 8$. Using Fintushel and Stern's technique in [FS3], J. Park, Stipsicz and Szabó were able to construct infinitely many irreducible non-symplectic smooth structures on $\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$ in [PSS]. The first *symplectic* exotic irreducible smooth structure on $\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$ was constructed in [A2] by the first author.

The existence of infinitely many irreducible smooth structures on $3\mathbb{CP}^2\#19\overline{\mathbb{CP}}^2$ was shown by Friedman, Morgan, O'Grady, Stipsicz and Szabó in [FM, MO, SS1]. Exotic smooth structures on $3\mathbb{CP}^2\#m\overline{\mathbb{CP}}^2$ for $14 \leq m \leq 18$ were first constructed by Gompf in [Go]. The existence of infinitely many irreducible smooth structures on $3\mathbb{CP}^2\#m\overline{\mathbb{CP}}^2$ for $14 \leq m \leq 18$ were shown by Stipsicz, Szabó and Yu in [St1, St2, Sz1, Yu]. Infinitely many irreducible smooth structures on $3\mathbb{CP}^2\#m\overline{\mathbb{CP}}^2$ for

---

Date: December 24, 2006. Revised on February 23, 2007.

2000 *Mathematics Subject Classification.* Primary 57R55; Secondary 57R17, 57M05.

$10 \leq m \leq 13$ were constructed by the second author in [P1, P2, P3]. Using rational blowdown techniques, Stipsicz and Szabó constructed infinitely many irreducible smooth structures on $3\mathbb{CP}^2\#9\overline{\mathbb{CP}}^2$ in [SS3], and J. Park constructed infinitely many irreducible smooth structures on $3\mathbb{CP}^2\#8\overline{\mathbb{CP}}^2$ in [Pa3]. The first exotic irreducible smooth structure on $3\mathbb{CP}^2\#7\overline{\mathbb{CP}}^2$ was constructed by the first author in [A2].

In this paper, we use the techniques and constructions in [A1] and [A2] to prove the following.

Theorem 1. *Let M be either $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$ or $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$. There exists a smooth closed simply-connected irreducible symplectic 4-manifold that is homeomorphic but not diffeomorphic to M.*

The main technical difficulty in the proof of Theorem 1 is the computation of the fundamental groups of our exotic 4-manifolds. Even though we cannot always completely pin down the fundamental groups at all stages of our construction, we are able to identify enough of the generators (up to conjugacy) and determine enough relations among them so that we are able to deduce, after careful choices of gluing, that the resulting 4-manifolds are simply-connected.

In the forthcoming paper [ABP] with R. İnanç Baykur, we will present some alternative constructions of an irreducible symplectic smooth structure on M. In [ABP], we also construct infinitely many irreducible non-symplectic smooth structures on M, $3\mathbb{CP}^2\#7\overline{\mathbb{CP}}^2$ and other small 4-manifolds.

Here is how our paper is organized. Section 2 contains some definitions and formulas that will be important throughout the paper. Section 3 quickly reviews the 4-manifolds that were constructed in [A1]. These 4-manifolds will then serve as some of the building blocks in constructing an exotic $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$ in Section 4 and an exotic $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$ in Section 5. Finally, Section 6 contains some technical computations which are crucial in showing that the fundamental groups of our exotic 4-manifolds are trivial.

2. Generalized fiber sum

We first present a few standard definitions that will be used throughout the paper.

Definition 2. Let X and Y be closed oriented smooth 4-manifolds each containing a smoothly embedded surface Σ of genus $g \geq 1$. Assume Σ represents a homology of infinite order and has self-intersection zero in X and Y, so that there exists a product tubular neighborhood, say $\nu\Sigma \cong \Sigma \times D^2$, in both X and Y. Using an orientation-reversing and fiber-preserving diffeomorphism $\psi : \Sigma \times S^1 \to \Sigma \times S^1$, we can glue $X \setminus \nu\Sigma$ and $Y \setminus \nu\Sigma$ along the boundary $\partial(\nu\Sigma) \cong \Sigma \times S^1$. The resulting closed oriented smooth 4-manifold, denoted $X\#_\psi Y$, is called a *generalized fiber sum* of X and Y along Σ.

Definition 3. Let $e(X)$ and $\sigma(X)$ denote the Euler characteristic and the signature of a closed oriented smooth 4-manifold X, respectively. We define

$$c_1^2(X) := 2e(X) + 3\sigma(X), \quad \chi_h(X) := \frac{e(X) + \sigma(X)}{4}.$$

If X is a complex surface, then $c_1^2(X)$ and $\chi_h(X)$ are the square of the first Chern class $c_1(X)$ and the holomorphic Euler characteristic, respectively. Note that these can be used as the coordinates for the "geography problem" for complex surfaces or irreducible smooth 4-manifolds (cf. [GS]).

For the simply-connected 4-manifolds $m\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2$, we have $e = 2+m+n$ and $\sigma = m-n$. Hence we get

$$c_1^2(m\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2) = 5m-n+4, \quad \chi_h(m\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2) = \frac{m+1}{2}.$$

Lemma 4. *Let X and Y be closed, oriented, smooth 4-manifolds containing an embedded surface Σ of self-intersection 0. Then*

$$\begin{aligned} c_1^2(X\#_\psi Y) &= c_1^2(X)+c_1^2(Y)+8(g-1), \\ \chi_h(X\#_\psi Y) &= \chi_h(X)+\chi_h(Y)+(g-1), \end{aligned}$$

where g is the genus of the surface Σ.

Proof. The above formulas simply follow from the well-known formulas

$$e(X\#_\psi Y) = e(X)+e(Y)-2e(\Sigma), \quad \sigma(X\#_\psi Y) = \sigma(X)+\sigma(Y). \qquad \square$$

If X and Y are symplectic 4-manifolds and Σ is a symplectic submanifold in both, then according to a theorem of Gompf (cf. [Go]), $X\#_\psi Y$ admits a symplectic structure. In such a case, we will call $X\#_\psi Y$ a *symplectic sum.*

3. Building blocks

We review the main construction in [A1]. Let K denote a trefoil knot in S^3. Let νK denote the tubular neighborhood of K in S^3. It is well-known (cf. [BZ]) that

$$\pi_1(S^3\setminus\nu K) = \langle a,b \mid aba=bab\rangle,$$

and b and ab^2ab^{-4} represent a meridian and a longitude of K, respectively. It is also well-known that $\gamma_1 = a^{-1}b$ and $\gamma_2 = b^{-1}aba^{-1}$ generate the image of the fundamental group of the genus one Seifert surface of K under the inclusion-induced homomorphism. Let M_K denote the result of 0-surgery on K. Clearly, we have

$$\begin{aligned} (1) \qquad \pi_1(M_K\times S^1) &= \pi_1(M_K)\oplus\mathbb{Z} \\ &= \langle a,b,x \mid aba=bab,\ ab^2ab^{-4}=1,\ [x,a]=[x,b]=1\rangle. \end{aligned}$$

Since K is a genus one fibered knot, M_K is a T^2 fiber bundle over S^1 with a section b. Thus $M_K\times S^1$ is a T^2 fiber bundle over T^2, with a section $S = b\times S^1 = b\times x$. From [Th], we know that there exists a symplectic form on $M_K\times S^1$ with respect to which both S and a torus fiber F are symplectic submanifolds. Let $\mathrm{pr}: M_K\times S^1\to S^1$ denote the projection map onto the second factor.

Lemma 5. *Let $C_S = (M_K\times S^1)\setminus\nu S$ be the complement of a tubular neighborhood of a section S in $M_K\times S^1$. Then we have*

$$\pi_1(C_S) = \langle a,b,x \mid aba=bab,\ [x,a]=[x,b]=1\rangle.$$

Proof. Note that $C_S = (M_K\setminus\nu b)\times S^1 = (S^3\setminus\nu K)\times S^1$. $\square$

Lemma 6. *Let $C_F = (M_K\times S^1)\setminus\nu F$ be the complement of a tubular neighborhood of a fiber F in $M_K\times S^1$. Then we have*

$$\begin{aligned} \pi_1(C_F) = \langle \gamma_1',\gamma_2',d,y \mid & [\gamma_1',\gamma_2'] = [y,\gamma_1'] = [y,\gamma_2'] = 1, \\ & d\gamma_1'd^{-1} = \gamma_1'\gamma_2',\ d\gamma_2'd^{-1} = (\gamma_1')^{-1}\rangle. \end{aligned}$$

Proof. C_F is homotopy equivalent to a T^2 fiber bundle over a wedge of two circles. The monodromy along the circle y is trivial whereas the monodromy along the circle d is the same as the monodromy of M_K. Recall that $\gamma_1' = c^{-1}d$ and $\gamma_2' = d^{-1}cdc^{-1}$. Thus we have

$$\begin{aligned} d\gamma_1'd^{-1} &= dc^{-1}dd^{-1} = dc^{-1} = \gamma_1'\gamma_2', \\ d\gamma_2'd^{-1} &= dd^{-1}cdc^{-1}d^{-1} = (cdc^{-1})d^{-1} \\ &= (d^{-1}cd)d^{-1} = d^{-1}c = (\gamma_1')^{-1}. \end{aligned}$$

Note that $cdc = dcd$ implies $cdc^{-1} = d^{-1}cd$. □

Lemma 7. *Let Y_K be the symplectic sum of two copies of $M_K \times S^1$, identifying a section S in one copy with a fiber F in the other copy. If the gluing map ψ satisfies $\psi_*(x) = \gamma_1'$ and $\psi_*(b) = \gamma_2'$, then*

$$\begin{aligned} \pi_1(Y_K) &= \langle a, b, x, \gamma_1', \gamma_2', d, y \mid aba = bab,\ [x,a] = [x,b] = 1, \\ &\quad [\gamma_1', \gamma_2'] = [y, \gamma_1'] = [y, \gamma_2'] = 1,\ d\gamma_1'd^{-1} = \gamma_1'\gamma_2',\ d\gamma_2'd^{-1} = (\gamma_1')^{-1}, \\ &\quad x = \gamma_1',\ b = \gamma_2',\ ab^2ab^{-4} = [d,y]\rangle \\ &= \langle a, b, x, d, y \mid aba = bab,\ [x,a] = [x,b] = 1, \\ &\quad [y,x] = [y,b] = 1,\ dxd^{-1} = xb,\ dbd^{-1} = x^{-1},\ ab^2ab^{-4} = [d,y]\rangle. \end{aligned}$$

Proof. By Seifert-Van Kampen Theorem, $\pi_1(Y_K) = (\pi_1(C_S) * \pi_1(C_F))/\pi_1(T^3)$. One circle factor of T^3 is identified with the longitude of K on one side and the meridian of the torus fiber in $M_K \times S^1$ on the other side. This gives the last relation. □

Inside Y_K, we can find a genus two symplectic submanifold Σ_2 which is the internal sum of a punctured fiber F_0 in C_S and a punctured section S_0 in C_F. The inclusion-induced homomorphism maps the standard generators of $\pi_1(\Sigma_2)$ to $a^{-1}b$, $b^{-1}aba^{-1}$, d and y in $\pi_1(Y_K)$.

Lemma 8. *There are nonnegative integers m and n such that*

$$\begin{aligned} \pi_1(Y_K \setminus \nu\Sigma_2) &= \langle a, b, x, d, y;\ g_1, \dots, g_m \mid aba = bab, \\ &\quad [y,x] = [y,b] = 1,\ dxd^{-1} = xb,\ dbd^{-1} = x^{-1}, \\ &\quad ab^2ab^{-4} = [d,y],\ r_1 = \cdots = r_n = 1,\ r_{n+1} = 1\rangle, \end{aligned} \tag{2}$$

where the generators $g_1, \dots, g_m$ and relators $r_1, \dots, r_n$ all lie in the normal subgroup N generated by the element $[x,b]$, and the relator r_{n+1} is a word in x, a and elements of N. Moreover, if we add an extra relation $[x,b] = 1$ to (2), then the relation $r_{n+1} = 1$ simplifies to $[x,a] = 1$.

Proof. This follows from Seifert-Van Kampen Theorem. Note that $[x,b]$ is a meridian of Σ_2 in Y_K. Hence setting $[x,b] = 1$ should turn $\pi_1(Y_K \setminus \nu\Sigma_2)$ into $\pi_1(Y_K)$. Also note that a is another meridian of K, and so $a \times S^1$ is another torus section of the fiber bundle $M_K \times S^1$. Thus $[x,a]$ is the boundary of a punctured torus section in $C_S \setminus \nu F_0$, and is no longer trivial in $\pi_1(Y_K \setminus \nu\Sigma_2)$. By setting $[x,b] = 1$, the relation $r_{n+1} = 1$ is to turn into $[x,a] = 1$.

It remains to check that the relations in $\pi_1(Y_K)$ other than $[x,a] = [x,b] = 1$ remain the same in $\pi_1(Y_K \setminus \nu\Sigma_2)$. By choosing a suitable point $\theta \in S^1$ away from $\mathrm{pr}|_{C_S}(F_0)$, the projection of the punctured fiber F_0 that forms a part of Σ_2, we obtain an embedding of the knot complement $(S^3 \setminus \nu K) \times \{\theta\} \hookrightarrow C_S \setminus \nu F_0$. This means that $aba = bab$ holds in $\pi_1(Y_K \setminus \nu\Sigma_2)$. Since $[\Sigma_2]^2 = 0$, there exists a parallel copy of Σ_2 outside $\nu\Sigma_2$, wherein the identity $ab^2ab^{-4} = [d, y]$ still holds. The other four remaining relations in $\pi_1(Y_K)$ are coming from the monodromy of a torus fiber bundle over a torus. Since these four relations will now describe the monodromy of a punctured torus fiber bundle over a punctured torus, they still hold true in $\pi_1(Y_K \setminus \nu\Sigma_2)$. □

Now take two copies of Y_K. Suppose that the fundamental group of the second copy has e, f, z, s, t as generators, and the inclusion-induced homomorphism in the second copy maps the generators of $\pi_1(\Sigma_2)$ to $e^{-1}f$, $f^{-1}efe^{-1}$, s and t in $\pi_1(Y_K)$. Let X_K denote the symplectic sum of two copies of Y_K along Σ_2, where the gluing map ψ maps the generators as follows:

$$\psi_*(a^{-1}b) = s,\ \psi_*(b^{-1}aba^{-1}) = t,\ \psi_*(d) = e^{-1}f,\ \psi_*(y) = f^{-1}efe^{-1}.$$

Lemma 9. *There are nonnegative integers m and n such that*

$$\begin{aligned}\pi_1(X_K) = \langle & a,b,x,d,y;\ e,f,z,s,t;\ g_1,\dots,g_m;\ h_1,\dots,h_m \mid \\ & aba = bab,\ [y,x] = [y,b] = 1, \\ & dxd^{-1} = xb,\ dbd^{-1} = x^{-1},\ ab^2ab^{-4} = [d,y], \\ & r_1 = \cdots = r_n = r_{n+1} = 1,\ r'_1 = \cdots = r'_n = r'_{n+1} = 1, \\ & efe = fef,\ [t,z] = [t,f] = 1, \\ & szs^{-1} = zf,\ sfs^{-1} = z^{-1},\ ef^2ef^{-4} = [s,t], \\ & d = e^{-1}f,\ y = f^{-1}efe^{-1},\ a^{-1}b = s,\ b^{-1}aba^{-1} = t, [x,b] = [z,f]\rangle,\end{aligned}$$

where g_i, h_i $(i = 1,\dots,m)$ and r_j, r'_j $(j = 1,\dots,n)$ all lie in the normal subgroup N generated by $[x,b] = [z,f]$. Moreover, r_{n+1} is a word in x, a and elements of N, and r'_{n+1} is a word in z, e and elements of N.

Proof. This is just a straightforward application of Seifert-Van Kampen Theorem and Lemma 8. □

Lemma 10. *There are nonnegative integers m, n, p and q such that*

$$\begin{aligned}\pi_1(X_K \setminus \nu\Sigma_2) = \langle & a,b,x,d,y;\ e,f,z,s,t;\ g_1,\dots,g_m;\ h_1,\dots,h_m;\ k_1,\dots,k_p \mid \\ & aba = bab,\ [y,x] = [y,b] = 1, \\ & dxd^{-1} = xb,\ dbd^{-1} = x^{-1},\ ab^2ab^{-4} = [d,y], \\ & r_1 = \cdots = r_n = r_{n+1} = 1,\ r'_1 = \cdots = r'_n = r'_{n+1} = 1, \\ & efe = fef,\ [t,z] = [t,f] = 1, \\ & szs^{-1} = zf,\ sfs^{-1} = z^{-1},\ ef^2ef^{-4} = [s,t], \\ & d = e^{-1}f,\ y = f^{-1}efe^{-1},\ a^{-1}b = s,\ b^{-1}aba^{-1} = t, \\ & r''_1 = \cdots = r''_q = 1\rangle,\end{aligned}$$

where the elements $g_1,\dots,g_m, r_1,\dots,r_n$ lie in the normal subgroup N generated by $[x,b]$, $h_1,\dots,h_m, r'_1,\dots,r'_n$ lie in the normal subgroup N' generated by $[z,f]$, and $k_1,\dots,k_p, r''_1\dots,r''_q$ lie in the normal subgroup N'' generated by $[x,b][z,f]^{-1}$. Moreover, r_{n+1} is a word in x, a and elements of N, and r'_{n+1} is a word in z, e and elements of N'.

Proof. Note that $[x,b][z,f]^{-1}$ is the meridian of Σ_2 in X_K. By Seifert-Van Kampen Theorem, setting $[x,b][z,f]^{-1} = 1$ turns $\pi_1(X_K \setminus \nu\Sigma_2)$ into $\pi_1(X_K)$. □

4. CONSTRUCTION OF AN EXOTIC $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$

In this section, we present a construction of a simply-connected, symplectic 4-manifold X that is homeomorphic but not diffemorphic to $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$. We will use the Seiberg-Witten invariant of X to distinguish it from $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$. Our first building block will be the symplectic 4-manifold X_K and a genus two symplectic submanifold $\Sigma_2 \subset X_K$ from Section 3. The other building block will be the 4-manifold $Y = T^4\#2\overline{\mathbb{CP}}^2$, the 4-torus blown up twice. X will then be the 4-manifold obtained by taking the symplectic sum of X_K and Y along Σ_2 and a certain genus two surface $\Sigma_2' \subset Y$.

We find a symplectically embedded genus two surface Σ_2' in Y as follows. First we introduce the notation $T_{i,j}$ $(1 \leq i < j \leq 4)$ for the 2-torus inside T^4 that has nontrivial ith and jth circle factors. For example, $T_{1,2} = S^1 \times S^1 \times \{\mathrm{pt}\} \times \{\mathrm{pt}\}$. Let $p_{i,j} : T^4 \to T_{i,j}$ be the projection map. Let $\omega_{i,j}$ be a standard product volume form on $T_{i,j}$.

Next we fix a factorization $T^4 = T^2 \times T^2$ and endow T^4 with a corresponding product symplectic form $\omega = p_{1,2}^*(\omega_{1,2}) + p_{3,4}^*(\omega_{3,4})$. Consider one copy of a horizontal torus $T_{1,2}$, and one copy of a vertical torus $T_{3,4}$. They are both symplectically embedded in T^4 with respect to ω. We symplectically resolve their intersection and obtain a symplectic surface of self-intersection 2. Next blow up twice to get a symplectic surface Σ_2' of self-intersection 0 in $Y = T^4\#2\overline{\mathbb{CP}}^2$.

Note that the fundamental group of Y is $\mathbb{Z}^4$. Let α_i $(i = 1, \dots, 4)$ denote the generators of $\pi_1(Y)$. The fundamental group of the complement of a tubular neighborhood $\nu\Sigma_2' \cong \Sigma_2' \times D^2$ is also $\mathbb{Z}^4$. It is also generated by the α_i's. This is because the normal circle $\mu' = \{\mathrm{pt}\} \times \partial D^2$ of $\nu\Sigma_2'$ (i.e. the meridian of Σ_2') can be deformed into one of the exceptional spheres, and thus is trivial in $\pi_1(Y \setminus \nu\Sigma_2')$.

Recall from Section 3 (cf. [A1]) that $a^{-1}b$, $b^{-1}aba^{-1}$, d, y, and $\mu = \{\mathrm{pt}\} \times S^1 = [x,b][z,f]^{-1}$ generate the inclusion-induced image of $\pi_1(\Sigma_2 \times S^1)$ inside $\pi_1(X_K \setminus \nu\Sigma_2)$. As before, let α_1, α_2, α_3, α_4 and μ' generate $\pi_1(\Sigma_2' \times S^1)$. We choose the gluing diffeomorphism $\psi : \Sigma_2 \times S^1 \to \Sigma_2' \times S^1$ that maps the fundamental group generators as follows:

$$\psi_*(a^{-1}b) = \alpha_1,\ \psi_*(b^{-1}aba^{-1}) = \alpha_2,\ \psi_*(d) = \alpha_3,\ \psi_*(y) = \alpha_4,\ \psi_*(\mu) = \mu'.$$

By Gompf's theorem in [Go], $X := X_K\#_\psi(T^4\#2\overline{\mathbb{CP}}^2)$ is symplectic.

Lemma 11. *X is simply-connected.*

Proof. By Seifert-Van Kampen Theorem, we have

$$\pi_1(X) = \frac{\pi_1(X_K \setminus \nu\Sigma_2) * \pi_1(Y \setminus \nu\Sigma_2')}{\langle a^{-1}b = \alpha_1,\ b^{-1}aba^{-1} = \alpha_2,\ d = \alpha_3,\ y = \alpha_4,\ [x,b] = [z,f]\rangle}.$$

Note that the normal circle $\mu = [x,b][z,f]^{-1}$ of $X_K \setminus \nu\Sigma_2$ becomes trivial in $\pi_1(X)$ since $\mu' = 1$. This implies that the generators $k_1, \dots, k_p$ of $\pi_1(X_K \setminus \nu\Sigma_2)$ are trivial and the relations $r_1'' = \cdots = r_q'' = 1$ are redundant in $\pi_1(X)$ (see Lemma 10). Since α_i's commute with one another, we get the following commutator relations in the fundamental group of X: $[a^{-1}b, b^{-1}aba^{-1}] = [a^{-1}b, d] = [a^{-1}b, y] = [b^{-1}aba^{-1}, d] = [b^{-1}aba^{-1}, y] = [d,y] = 1$. In summary we get the following presentation for the fundamental group of X.

$$\begin{aligned}\pi_1(X) = \langle & a,b,x,d,y;\ e,f,z,s,t;\ g_1,\dots,g_m;\ h_1,\dots,h_m \mid \\ & aba=bab,\ [y,x]=[y,b]=1, \\ & dxd^{-1}=xb,\ dbd^{-1}=x^{-1},\ ab^2ab^{-4}=[d,y], \\ & r_1=\dots=r_n=r_{n+1}=1,\ r_1'=\dots=r_n'=r_{n+1}'=1, \\ & efe=fef,\ [t,z]=[t,f]=1, \\ & szs^{-1}=zf,\ sfs^{-1}=z^{-1},\ ef^2ef^{-4}=[s,t], \\ & d=e^{-1}f,\ y=f^{-1}efe^{-1},\ a^{-1}b=s,\ b^{-1}aba^{-1}=t, \\ & [s,t]=[s,d]=[s,y]=[t,d]=[t,y]=[d,y]=1\rangle.\end{aligned} \tag{3}$$

To prove $\pi_1(X)=1$, it is enough to prove that $b=d=f=s=1$, since these will imply that all other generators are trivial. It is not hard to show that the following five identities hold in $\pi_1(X)$. The proof of these identities is postponed to the Appendix (see Section 6).

$$dbd = bd^2b, \tag{4}$$

$$bsb = sb^2s, \tag{5}$$

$$sfs = fs^2f, \tag{6}$$

$$fdf = df^2d, \tag{7}$$

$$(bdb^{-1})s = s(bdb^{-1}). \tag{8}$$

Now rewrite (4) as $(d^{-1}b^{-1}d)(bdb^{-1})=d$. Since s commutes with d and bdb^{-1}, we conclude that s also commutes with $d^{-1}b^{-1}d$. It follows that s commutes with the inverse of $d^{-1}b^{-1}d$, i.e. $s(d^{-1}bd)=(d^{-1}bd)s$. Since s commutes with d, we must have $sb=bs$. Using $sb=bs$ in (5), we get $s=1$. Then (6) implies that $f=1$. Similarly, (7) and (4) in turn imply that $d=b=1$. Thus $\pi_1(X)$ is trivial. □

From Lemma 10, it is easy to see that the abelianization of $\pi_1(X_K\setminus\nu\Sigma_2)$ is trivial. By suitably adjoining an abelian group $\pi_1(Y\setminus\nu\Sigma_2')\cong\mathbb{Z}^4$ to $\pi_1(X_K\setminus\nu\Sigma_2)$, we were able to infuse enough commutativity to eventually eliminate all the generators in the above proof. Note that the relations $ab^2ab^{-4}=[d,y]$ and $ef^2ef^{-4}=[s,t]$ in (3) are not used in the above proof or in the proof of (4)–(8) in the Appendix. An astute reader will notice that this and the fact that the other meridians are eventually set to 1 allow us to be a little lax on the orientation convention for meridians.

Lemma 12. *$e(X)=10$, $\sigma(X)=-2$, $c_1^2(X)=14$, and $\chi_h(X)=2$.*

Proof. Let $Y=T^4\#2\overline{\mathbb{CP}}^2$ as before. We know that $e(X)=e(X_K)+e(Y)+4$, $\sigma(X)=\sigma(X_K)+\sigma(Y)$, $c_1^2(X)=c_1^2(X_K)+c_1^2(Y)+8$, and $\chi_h(X)=\chi_h(X_K)+\chi_h(Y)+1$. We easily compute that $e(Y)=2$, $\sigma(Y)=-2$, $c_1^2(Y)=-2$, and $\chi_h(Y)=0$. Since $e(X_K)=4$, $\sigma(X_K)=0$, $c_1^2(X_K)=8$ and $\chi_h(X_K)=1$ (cf. [A1]), our results follow. □

From Freedman's classification theorem (cf. [Fre]) for simply-connected topological 4-manifolds, we conclude that X is homeomorphic to $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$. It follows from Taubes's theorem (cf. [Ta]) that $\mathrm{SW}_X(K_X)=\pm1$, where K_X is the canonical class of X. Next we apply the connected sum theorem (cf. [Wi]) for the Seiberg-Witten invariant to deduce that the Seiberg-Witten invariant is trivial for

$3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$. Since the Seiberg-Witten invariant is a diffeomorphism invariant, we conclude that X is not diffeomorphic to $3\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$.

Since $\pi_2(M_K \times S^1) = 0$, we conclude that $M_K \times S^1$ is a minimal symplectic 4-manifold. Usher's theorem (cf. [Us]) then implies that the symplectic sums Y_K and X_K are minimal as well. Both exceptional spheres E_1 and E_2 in Y transversely intersect the genus two surface Σ_2' once since $[\Sigma_2'] = [T_{1,2}] + [T_{3,4}] - [E_1] - [E_2] \in H_2(Y;\mathbb{Z})$. Usher's theorem once again implies that the symplectic sum X is a minimal symplectic 4-manifold. Since symplectic minimality implies irreducibility for simply-connected 4-manifolds (cf. [Ko2] for $b_2^+ > 1$ case), X is also smoothly irreducible.

5. Construction of an exotic $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$

In this section, we construct a simply-connected symplectic 4-manifold U homeomorphic but not diffemorphic to $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$. Using Usher's Theorem (cf. [Us]), we will distinguish U from $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$.

The manifold U will be the symplectic sum of the 4-manifold Y_K in Section 3 and $Q = (M_K \times S^1)\#2\overline{\mathbb{CP}}^2$ along the genus two surfaces Σ_2 and Σ_2''. The symplectic genus two submanifold $\Sigma_2'' \subset Q$ is obtained by symplectically resolving the intersection of a torus fiber F and a torus section S of $M_K \times S^1$ (see Section 3) and then blowing up at two points.

Let g, h, z be generators of $\pi_1(M_K \times S^1)$ as in (1) such that the inclusion-induced homomorphisms map the generators of $\pi_1(F)$ to $g^{-1}h$ and $h^{-1}ghg^{-1}$, and the generators of $\pi_1(S)$ to h and z. Let us choose the gluing diffeomorphism $\varphi : \Sigma_2 \times S^1 \to \Sigma_2'' \times S^1$ that maps the elements $a^{-1}b$, $b^{-1}aba^{-1}$, d, y and μ of $\pi_1(Y_K \setminus \nu\Sigma_2)$ to the elements $g^{-1}h$, $h^{-1}ghg^{-1}$, h, z and μ'' of $\pi_1(Q \setminus \nu\Sigma_2'')$ according to the following rule:

$$\varphi_*(a^{-1}b) = h,\ \varphi_*(b^{-1}aba^{-1}) = z,\ \varphi_*(d) = g^{-1}h,\ \varphi_*(y) = h^{-1}ghg^{-1},\ \varphi_*(\mu) = \mu''.$$

Here, μ and μ'' denote the meridians of Σ and Σ_2''. It follows from Gompf's theorem (cf. [Go]) that $U := Y_K\#_\varphi((M_K \times S^1)\#2\overline{\mathbb{CP}}^2)$ is symplectic.

Lemma 13. *U is simply-connected.*

Proof. By Seifert-Van Kampen Theorem, we have

$$\pi_1(U) = \frac{\pi_1(Y_K \setminus \nu\Sigma_2) * \pi_1(Q \setminus \nu\Sigma_2'')}{\langle a^{-1}b = h,\ b^{-1}aba^{-1} = z,\ d = g^{-1}h,\ y = h^{-1}ghg^{-1},\ \mu = 1\rangle}.$$

Since μ'' is nullhomotopic in $Q \setminus \nu\Sigma_2''$, the normal circle $\mu = [x, b]$ of $\pi_1(Y_K \setminus \nu\Sigma_2)$ becomes trivial in $\pi_1(U)$. This in turn implies that the generators $g_1, \dots, g_m$ of $\pi_1(Y_K \setminus \nu\Sigma_2)$ become trivial in $\pi_1(U)$, the relations $r_1 = \cdots = r_n = 1$ become redundant, and the relation $r_{n+1} = 1$ turns into $[x, a] = 1$ (see Lemma 8). Since $\pi_1(Q \setminus \nu\Sigma_2'') \cong \pi_1(Q) \cong \pi_1(M_K \times S^1)$, we conclude that $\pi_1(U)$ is generated by

$$a,\ b,\ x,\ d,\ y;\ g,\ h,\ z.$$

The following relations hold in $\pi_1(U)$.

$$\begin{aligned} & aba = bab,\ [x, b] = [x, a] = [y, x] = [y, b] = 1, \\ & dxd^{-1} = xb,\ dbd^{-1} = x^{-1},\ ab^2ab^{-4} = [d, y], \\ & ghg = hgh,\ gh^2gh^{-4} = 1,\ [z, g] = [z, h] = 1, \\ & a^{-1}b = h,\ b^{-1}aba^{-1} = z,\ d = g^{-1}h,\ y = h^{-1}ghg^{-1}. \end{aligned} \tag{9}$$

Using the above relations, we can obtain the following six identities in $\pi_1(U)$. The proof of these identities is postponed to the Appendix (see Section 6).

$$bhb = hb^2h \tag{10}$$

$$d(hbh^{-1}) = (hbh^{-1})d \tag{11}$$

$$h(bdb^{-1}) = (bdb^{-1})h \tag{12}$$

$$d^2 = b^{-1}dbdb^{-1} \tag{13}$$

$$b^2 = hbh^{-1}bh \tag{14}$$

$$h^2 = dhd^{-1}hd \tag{15}$$

Using (10), we have $hbh^{-1} = b^{-1}hb^2$. Notice that by (11) hbh^{-1} commutes with d. Then d commutes with $b^{-1}hb^2$, so we have $d(b^{-1}hb^2) = (b^{-1}hb^2)d$. This in turn implies that $(bdb^{-1})hb^2 = hb^2d$. Since h commutes with bdb^{-1} by (12), we have $hbdb^{-1}b^2 = hb^2d$. The last equality implies that $db = bd$. Using this fact in (13), (14) and (15), we get $b = h = d = 1$. Now it easily follows from the relations of $\pi_1(U)$ that $a = x = y = g = z = 1$. Thus we have proved that $\pi_1(U)$ is trivial. □

Lemma 14. *$e(U) = 6$, $\sigma(U) = -2$, ${c_1}^2(U) = 6$, and $\chi_h(U) = 1$*

Proof. Let $Q = (M_K \times S^1)\#2\overline{\mathbb{CP}}^2$. We have $e(U) = e(Y_K) + e(Q) + 4$, $\sigma(U) = \sigma(Y_K)+\sigma(Q)$, $c_1^2(U) = c_1^2(Y_K)+c_1^2(Q)+8$, and $\chi_h(U) = \chi_h(Y_K)+\chi_h(Q)+1$. Since $e(Q) = 2$, $\sigma(Q) = -2$, $c_1^2(Q) = -2$, $\chi_h(Q) = 0$, and $e(Y_K) = \sigma(Y_K) = c_1^2(Y_K) = \chi_h(Y_K) = 0$ (cf. [A1]), our results follow. □

By Freedman's theorem (cf. [Fre]) and the lemmas above, we deduce that U is homeomorphic to $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$. Note that U is a symplectic sum of a non-minimal 4-manifold $Q = (M_K \times S^1)\#2\overline{\mathbb{CP}}^2$ with a minimal 4-manifold Y_K. Both exceptional spheres E_1 and E_2 in Q transversely intersect the genus two surface Σ_2'' once since we have $[\Sigma_2''] = [F] + [S] - [E_1] - [E_2] \in H_2(Q;\mathbb{Z})$. It follows from Usher's Theorem (cf. [Us]) that U is a minimal symplectic 4-manifold. Since symplectic minimality implies irreducibility for simply-connected 4-manifolds (cf. [HK] for $b_2^+ = 1$ case), U is also smoothly irreducible. We conclude that it cannot be diffeomorphic to $\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$.

6. Appendix

In this section we fill in the details left out in the proofs of Lemmas 11 and 13 by proving the identities (4)–(8) in $\pi(X)$ and (10)–(15) in $\pi_1(U)$.

Proof of (4)–(8). From presentation (3), we get $x = db^{-1}d^{-1}$. Substituting this into the relation $dxd^{-1} = xb$, we get $d^2b^{-1}d^{-2} = db^{-1}d^{-1}b$, which can be rearranged to $b^{-1}d^{-2}b^{-1} = d^{-1}b^{-1}d^{-1}$. By taking inverses of both sides, we get (4). Similarly, substituting $z = sf^{-1}s^{-1}$ into the relation $szs^{-1} = zf$ yields $s^2f^{-1}s^{-2} = sf^{-1}s^{-1}f$. Rearrange this into $f^{-1}s^{-2}f^{-1} = s^{-1}f^{-1}s^{-1}$ and take inverses to obtain (6).

Next note that $a = bs^{-1}$, and thus $aba = bab$ implies that $bs^{-1}b^2s^{-1} = b^2s^{-1}b$. Hence $b^{-1}s^{-1}b^2s^{-1}b^{-1} = s^{-1}$, and by taking the inverses of both sides, we obtain

$$(bsb^{-1})(b^{-1}sb) = s. \tag{16}$$

Now $t = b^{-1}aba^{-1} = s^{-1}bsb^{-1}$. Since $st = ts$, we have $bsb^{-1} = s^{-1}bsb^{-1}s$. From (16), we get $b^2sb^{-2}s = bsb^{-1}$. Combining the last two identities, we obtain

$b^2sb^{-2}s = s^{-1}bsb^{-1}s$, which simplifies to $b^2sb^{-1} = s^{-1}bs$. The last identity is easily changed to (5).

Similarly, note that $e = fd^{-1}$, and thus $efe = fef$ implies that $fd^{-1}f^2d^{-1} = f^2d^{-1}f$. Then $f^{-1}d^{-1}f^2d^{-1}f^{-1} = d^{-1}$, and so

$$fdf^{-2}df = d. \tag{17}$$

Now $y = f^{-1}efe^{-1} = d^{-1}fdf^{-1}$. Since $dy = yd$, we have $fdf^{-1} = d^{-1}fdf^{-1}d$. From (17), we get $f^2df^{-2}d = fdf^{-1}$. Combining the last two identities, we obtain $f^2df^{-2}d = d^{-1}fdf^{-1}d$, which simplifies to $f^2df^{-1} = d^{-1}fd$. The last identity is easily changed to (7).

Finally, since $dt = td$, we have $ds^{-1}bsb^{-1} = s^{-1}bsb^{-1}d$. Since d commutes with s and s^{-1}, we conclude that

$$d(bsb^{-1}) = (bsb^{-1})d.$$

From (16), we also know that d commutes with the product $(bsb^{-1})(b^{-1}sb)$. It follows that d commutes with $b^{-1}sb$, i.e.

$$d(b^{-1}sb) = (b^{-1}sb)d. \tag{18}$$

Now it is easy to see that (18) can be rearranged to (8).

Proof of (10)–(15). From (9), $x = db^{-1}d^{-1}$. Plugging this into $dxd^{-1} = xb$, we get $d^2b^{-1}d^{-2} = db^{-1}d^{-1}b$. This implies that $d^{-2} = bd^{-1}b^{-1}d^{-1}b$, which gives (13).

Next we have $a^{-1}b = h$, and hence $a = bh^{-1}$. Plugging this last identity into $aba = bab$, we get $bh^{-1}b^2h^{-1} = b^2h^{-1}b$. This simplifies to $h^{-1}b^2h^{-1} = bh^{-1}b$, which can be easily rearranged to (14).

Similarly, we have $d = g^{-1}h$, and hence $g = hd^{-1}$. Plugging the last identity into $ghg = hgh$, we get $hd^{-1}h^2d^{-1} = h^2d^{-1}h$. It follows that $d^{-1}h^2d^{-1} = hd^{-1}h$, which can be easily arranged to (15).

The relation $h = a^{-1}b$ gives $a = bh^{-1}$. Thus $z = b^{-1}aba^{-1} = h^{-1}bhb^{-1}$. Since $[z, h] = 1$, we get $h^{-1}bhb^{-1}h = bhb^{-1}$. This simplifies to

$$(bhb^{-1})h = h(bhb^{-1}). \tag{19}$$

From (14), we have $b^2 = hbh^{-1}bh$, which can be rearranged as $hb^{-1}h^{-1}b = bhb^{-1}$. Using (19), we have $bhb^{-1} = h^{-1}(bhb^{-1})h = b^{-1}h^{-1}bh$. The identity $bhb^{-1} = b^{-1}h^{-1}bh$ can be easily rearranged to (10).

The relation $g^{-1}h = d$ gives $g = hd^{-1}$. Thus $y = h^{-1}ghg^{-1} = d^{-1}hdh^{-1}$. Since $[y, b] = 1$, we conclude that

$$d^{-1}(hdh^{-1})b = bd^{-1}(hdh^{-1}). \tag{20}$$

Now (15) implies that $hdh^{-1} = dh^{-1}d^{-1}h$. Substituting this last identity into (20), we obtain $h^{-1}d^{-1}hb = bh^{-1}d^{-1}h$. This implies that $d^{-1}(hbh^{-1}) = (hbh^{-1})d^{-1}$, which can be rearranged to (11).

Finally, note that z commutes with both g and h, so it must commute with $d = g^{-1}h$. In the proof of (10) we saw that $z = h^{-1}bhb^{-1}$. Substitute this into $dz = zd$ and obtain

$$d(h^{-1}bhb^{-1}) = (h^{-1}bhb^{-1})d. \tag{21}$$

From (14), we have $h^{-1}bh = b^{-1}h^{-1}b^2$. Substituting this into (21), we obtain $db^{-1}h^{-1}b = b^{-1}h^{-1}bd$. This can be rearranged to $(bdb^{-1})h^{-1} = h^{-1}(bdb^{-1})$, which implies (12).

ACKNOWLEDGMENTS

The authors thank R. İnanç Baykur and Ronald J. Stern for helpful discussions. Some of the computations have been double-checked by using the computer software package GAP (cf. [GAP]). A. Akhmedov was partially supported by the NSF grant FRG-0244663. B. D. Park was partially supported by CFI, NSERC and OIT grants.

REFERENCES

[A1] A. Akhmedov: Construction of new symplectic cohomology $S^2 \times S^2$, math.GT/0611128.

[A2] A. Akhmedov: Small exotic 4-manifolds, math.GT/0612130.

[ABP] A. Akhmedov, R. İ. Baykur and B. D. Park: Constructing infinitely many smooth structures on small 4-manifolds, preprint.

[Ba] R. Barlow: A simply connected surface of general type with $p_g = 0$, *Invent. Math.* **79** (1985), 293–301.

[BZ] G. Burde and H. Zieschang: *Knots.* Second edition, de Gruyter Studies in Math., 5, Walter de Gruyter & Co., Berlin, 2003.

[Do] S. K. Donaldson: Irrationality and the h-cobordism conjecture, *J. Differential Geom.* **26** (1987), 141–168.

[FS1] R. Fintushel and R. J. Stern: Rational blowdowns of smooth 4-manifolds, *J. Differential Geom.* **46** (1997), 181–235.

[FS2] R. Fintushel and R. J. Stern: Knots, links and 4-manifolds, *Invent. Math.* **134** (1998), 363–400.

[FS3] R. Fintushel and R. J. Stern: Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$, *J. Amer. Math. Soc.* **19** (2006), 171–180.

[Fre] M. H. Freedman: The topology of four-dimensional manifolds, *J. Differential Geom.* **17** (1982), 357–453.

[Fri] R. Friedman: Vector bundles and SO(3)-invariants for elliptic surfaces, *J. Amer. Math. Soc.* **8** (1995), 29–139.

[FM] R. Friedman and J. W. Morgan: *Smooth Four-Manifolds and Complex Surfaces.* Ergeb. Math. Grenzgeb. (3), 27, Springer-Verlag, Berlin, 1994.

[GAP] The GAP Group: GAP – Groups, Algorithms, and Programming, Version 4.3; 2002. (`http://www.gap-system.org`)

[Go] R. E. Gompf: A new construction of symplectic manifolds, *Ann. of Math.* **142** (1995), 527–595.

[GS] R. E. Gompf and A. I. Stipsicz: 4-*Manifolds and Kirby Calculus.* Grad. Stud. Math., 20, Amer. Math. Soc., Providence, RI, 1999.

[HK] M. J. D. Hamilton and D. Kotschick: Minimality and irreducibility of symplectic four-manifolds, *Int. Math. Res. Not.* **2006**, Art. ID 35032, 13 pp.

[Ko1] D. Kotschick: On manifolds homeomorphic to $\mathbb{CP}^2\#8\overline{\mathbb{CP}}^2$, *Invent. Math.* **95** (1989), 591–600.

[Ko2] D. Kotschick: The Seiberg-Witten invariants of symplectic four-manifolds (after C. H. Taubes), Séminaire Bourbaki, 1995/96, no. 812, *Astérisque* **241** (1997), 195–220.

[MO] J. W. Morgan and K. G. O'Grady: *Differential Topology of Complex Surfaces. Elliptic Surfaces with* $p_g = 1$ *: Smooth Classification.* Lecture Notes in Math., 1545, Springer-Verlag, Berlin, 1993.

[P1] B. D. Park: Exotic smooth structures on $3\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2$, *Proc. Amer. Math. Soc.* **128** (2000), 3057–3065.

[P2] B. D. Park: Exotic smooth structures on $3\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2$, Part II, *Proc. Amer. Math. Soc.* **128** (2000), 3067–3073.

[P3] B. D. Park: Constructing infinitely many smooth structures on $3\mathbb{CP}^2\#n\overline{\mathbb{CP}}^2$, *Math. Ann.* **322** (2002), 267–278.

[Pa1] J. Park: Seiberg-Witten invariants of generalised rational blow-downs, *Bull. Austral. Math. Soc.* **56** (1997), 363–384.

[Pa2] J. Park: Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$, *Invent. Math.* **159** (2005), 657–667.

[Pa3] J. Park: Exotic smooth structures on $3\mathbb{CP}^2\#8\overline{\mathbb{CP}}^2$, *Bull. London Math. Soc.* (to appear).

[PSS] J. Park, A. I. Stipsicz, and Z. Szabó: Exotic smooth structures on $\mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$, *Math. Res. Lett.* **12** (2005), 701–712.

[St1] A. I. Stipsicz: Donaldson invariants of certain symplectic manifolds, *J. Reine Angew. Math.* **465** (1995), 1-10.

[St2] A. I. Stipsicz: Donaldson series and (-1)-tori, *J. Reine Angew. Math.* **465** (1995), 133–144.

[SS1] A. I. Stipsicz and Z. Szabó: The smooth classification of elliptic surfaces with $b^+ > 1$, *Duke Math. J.* **75** (1994), 1–50.

[SS2] A. I. Stipsicz and Z. Szabó: An exotic smooth structure on $\mathbb{CP}^2\#6\overline{\mathbb{CP}}^2$, *Geom. Topol.* **9** (2005), 813–832.

[SS3] A. I. Stipsicz and Z. Szabó: Small exotic 4-manifolds with $b_2^+ = 3$, *Bull. London Math. Soc.* **38** (2006), 501–506.

[Sz1] Z. Szabó: Irreducible four-manifolds with small Euler characteristics, *Topology* **35** (1996), 411–426.

[Sz2] Z. Szabó: Exotic 4-manifolds with $b_2^+ = 1$, *Math. Res. Lett.* **3** (1996), 731–741.

[Ta] C. H. Taubes: The Seiberg-Witten invariants and symplectic forms, *Math. Res. Lett.* **1** (1994), 809–822.

[Th] W. P. Thurston: Some simple examples of symplectic manifolds, *Proc. Amer. Math. Soc.* **55** (1976), 467–468.

[Us] M. Usher: Minimality and symplectic sums, *Internat. Math. Res. Not.* **2006**, Art. ID 49857, 17 pp.

[Wi] E. Witten: Monopoles and four-manifolds, *Math. Res. Lett.* **1** (1994), 769–796.

[Yu] B. Z. Yu: Some computations of Donaldson's polynomials via flat connections, Ph. D. Thesis, U. C. Berkeley, 1994.

School of Mathematics, Georgia Institute of Technology, Atlanta, GA, 30332-0160, USA
E-mail address: `ahmadov@math.gatech.edu`

Department of Pure Mathematics, University of Waterloo, Waterloo, ON, N2L 3G1, Canada
E-mail address: `bdpark@math.uwaterloo.ca`